

Appendix:

CURRENT AS Registry Investigators

List of Investigators

Principal Investigators

Takeshi Kimura, Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

Ryuzo Sakata, Department of Cardiovascular Surgery, Kyoto University Graduate School of Medicine, Kyoto, Japan

List of participating centers and investigators for the CURRENT AS registry Cardiology

Kyoto University Graduate School of Medicine: Takeshi Kimura, Tomohiko Taniguchi, Hiroki Shiomi, Naritatsu Saito, Masao Imai, Junichi Tazaki, Toshiaki Toyota, Hirooki Higami, Tetsuma Kawaji

Kokura Memorial Hospital: Kenji Ando, Shinichi Shirai, Kengo Kourai, Takeshi Arita, Shiro Miura, Kyohei Yamaji

Shimada Municipal Hospital: Takeshi Aoyama, Norio Kanamori

Shizuoka City Shizuoka Hospital: Tomoya Onodera, Koichiro Murata

Kobe City Medical Center General Hospital: Yutaka Furukawa, Takeshi Kitai, Kitae Kim

Kurashiki Central Hospital: Kazushige Kadota, Yuichi Kawase, Keiichiro Iwasaki, Hiroshi Miyawaki, Ayumi Misao, Akimune Kuwayama, Masanobu Ohya, Takenobu Shimada, Hidewo Amano **Tenri Hospital:** Yoshihisa Nakagawa, Chisato Izumi, Makoto Miyake, Masashi Amano, Yusuke

Takahashi, Yusuke Yoshikawa, Shunsuke Nishimura, Maiko Kuroda

Nara Hospital, Kinki University Faculty of Medicine: Manabu Shirotani, Hirokazu Mitsuoka

Mitsubishi Kyoto Hospital: Shinji Miki, Tetsu Mizoguchi, Masashi Kato, Takafumi Yokomatsu, Akihiro Kushiyama, Hidenori Yaku, Toshimitsu Watanabe

Kinki University Hospital: Shunichi Miyazaki, Yutaka Hirano

Kishiwada City Hospital: Mitsuo Matsuda, Shintaro Matsuda, Sachiko Sugioka

Osaka Red Cross Hospital: Tsukasa Inada, Kazuya Nagao, Naoki Takahashi, Kohei Fukuchi

Koto Memorial Hospital: Tomoyuki Murakami, Hiroshi Mabuchi, Teruki Takeda, Tomoko Sakaguchi, Keiko Maeda, Masayuki Yamaji, Motoyoshi Maenaka, Yutaka Tadano

Shizuoka General Hospital: Hiroki Sakamoto, Yasuyo Takeuchi, Makoto Motooka, Ryusuke Nishikawa

Nishikobe Medical Center: Hiroshi Eizawa, Keiichiro Yamane, Mitsunori Kawato, Minako Kinoshita, Kenji Aida

Japanese Red Cross Wakayama Medical Center: Takashi Tamura, Mamoru Toyofuku, Kousuke

Takahashi, Euihong Ko

National Hospital Organization Kyoto Medical Center: Masaharu Akao, Mitsuru Ishii, Nobutoyo Masunaga, Hisashi Ogawa, Moritake Iguchi, Takashi Unoki, Kensuke Takabayashi, Yasuhiro Hamatani, Yugo Yamashita

The Tazuke Kofukai Medical Research Institute, Kitano Hospital: Moriaki Inoko, Eri Minamino-Muta, Takao Kato

Hikone Municipal Hospital: Yoshihiro Himura, Tomoyuki Ikeda Kansai Electric Power Hospital: Katsuhisa Ishii, Akihiro Komasa

Hyogo Prefectural Amagasaki General Medical Center: Yukihito Sato, Kozo Hotta, Shuhei Tsuji

Rakuwakai Otowa Hospital: Yuji Hiraoka, Nobuya Higashitani

Saiseikai Noe Hospital: Ichiro Kouchi, Yoshihiro Kato

Shiga Medical Center for Adults: Shigeru Ikeguchi, Yasutaka Inuzuka, Soji Nishio, Jyunya Seki

Hamamatsu Rosai Hospital: Eiji Shinoda, Miho Yamada, Akira Kawamoto, Chiyo Maeda

Japanese Red Cross Otsu Hospital: Takashi Konishi, Toshikazu Jinnai, Kouji Sogabe, Michiya

Tachiiri, Yukiko Matsumura, Chihiro Ota

Hirakata Kohsai Hospital: Shoji Kitaguchi, Yuko Morikami

Cardiovascular Surgery

Kyoto University Graduate School of Medicine: Ryuzo Sakata, Kenji Minakata, Kenji Minatoya

Kokura Memorial Hospital: Michiya Hanyu

Shizuoka City Shizuoka Hospital: Fumio Yamazaki

Kobe City Medical Center General Hospital: Tadaaki Koyama

Kurashiki Central Hospital: Tatsuhiko Komiya

Tenri Hospital: Kazuo Yamanaka

Nara Hospital, Kinki University Faculty of Medicine: Noboru Nishiwaki

Mitsubishi Kyoto Hospital: Hiroyuki Nakajima, Motoaki Ohnaka, Hiroaki Osada, Katsuaki Meshii

Kinki University Hospital: Toshihiko Saga Kishiwada City Hospital: Masahiko Onoe Osaka Red Cross Hospital: Shogo Nakayama Shizuoka General Hospital: Genichi Sakaguchi

Japanese Red Cross Wakayama Medical Center: Atsushi Iwakura

National Hospital Organization Kyoto Medical Center: Kotaro Shiraga

The Tazuke Kofukai Medical Research Institute, Kitano Hospital: Koji Ueyama

Hyogo Prefectural Amagasaki General Medical Center: Keiichi Fujiwara

Rakuwakai Otowa Hospital: Atsushi Fukumoto Shiga Medical Center for Adults: Masaki Park Hamamatsu Rosai Hospital: Junichiro Nishizawa Japanese Red Cross Otsu Hospital: Mitsuru Kitano

A clinical event committee

Hirotoshi Watanabe, MD (Kyoto University Graduate School of Medicine); Kenji Nakatsuma, MD (Kyoto University Graduate School of Medicine), Tomoki Sasa, MD (Kishiwada City Hospital)

Data S1.

Definitions of the clinical events

Death was regarded as cardiovascular in origin unless obvious non-cardiovascular causes could be identified. Sudden death was defined as unexplained death in previously stable patients. Any death during the hospitalization for aortic valve replacement or transcatheter aortic valve implantation was regarded as aortic valve procedure-related death. Aortic valve-related death included aortic valve procedure-related death, sudden death, and death due to heart failure related to aortic stenosis. Heart failure hospitalization was defined as hospitalization due to worsening heart failure requiring intravenous drug therapy.


Table S1. Clinical factors associated with sudden death in the non-HD cohort in the univariate and multivariable analyses.

Variables	Unadjusted HR* (95% CI)	P value	Adjusted HR (95% CI)	P value
Prior myocardial infarction	2.95 (1.90-4.43)	< 0.001	3.25 (1.59-6.61)	0.001
BMI <22	1.87 (1.26-2.85)	0.002	1.53 (0.98-2.41)	0.06
LVEF < 60%	2.71 (1.90-3.85)	< 0.001	1.56 (1.04-2.35)	0.03
Vmax ≥5m/s	1.67 (1.002-2.63)	0.049	1.50 (0.91-2.49)	0.11
Symptoms related to AS	2.34 (1.65-3.36)	< 0.001	1.40 (0.93-2.12)	0.11
Age ≥80	2.13 (1.47-3.12)	< 0.001	1.19 (0.78-1.82)	0.42
Male	1.52 (1.06-2.16)	0.02	1.53 (1.04-2.24)	0.03
TRPG ≥40 mmHg	1.71 (1.09-2.60)	0.02	1.16 (0.69-1.94)	0.58
Any valvular disease (moderate-severe)	1.37 (0.96-1.94)	0.08	1.01 (0.67-1.52)	0.98
Prior symptomatic stroke	1.18 (0.71-1.86)	0.50	0.95 (0.58-1.56)	0.84
Atrial fibrillation or flutter	1.07 (0.70-1.59)	0.75	1.01 (0.66-1.55)	0.95
Chronic lung disease (moderate-severe)	1.18 (0.42-2.60)	0.72	0.82 (0.33-1.99)	0.65
Malignancy currently under treatment	1.47 (0.57-3.07)	0.38	1.01 (0.44-2.36)	0.98
Aortic/peripheral disease	1.06 (0.45-2.12)	0.88	0.79 (0.35-1.74)	0.55
Current smoking	0.84 (0.33-1.75)	0.67	0.94 (0.41-2.19)	0.89
Diabetes on insulin	1.83 (0.86-3.39)	0.11	1.26 (0.60-2.63)	0.54
Coronary artery disease	1.47 (0.998-2.12)	0.052	0.60 (0.33-1.11)	0.10
Anemia	1.87 (1.31-2.69)	< 0.001	0.96 (0.65-1.43)	0.85
Hypertension	1.51 (1.004-2.33)	0.048	1.45 (0.94-2.24)	0.09

^{*} Competing risk of death other than sudden death was not taken into account in the univariate analysis, while it was taken into account in the multivariable analysis. Patients who had AVR or TAVI were censored at AVR/TAVI.

CI=confidence interval; HR=hazard ratio; TAVI=transcatheter aortic valve implantation. Other abbreviations were same as in Table 1.

Figure S1. Cumulative incidence of sudden cardiac death in HD and non-HD patients with severe AS.



Cumulative incidence of sudden death was estimated by the Gray method, taking the competing risk of death other than sudden death. AVR=aortic valve replacement; HD=hemodialysis;

TAVI=transcatheter aortic valve implantation.